

TYPE OF VACCINE FOR REVACCINATION

Revaccination against CVI is recommended to be conducted by types of vaccines (inactivated vaccine, vector vaccine, m-RNA vaccine) that were not used during the initial course of vaccination against CVI.

Revaccination against CVI is allowed to be conducted with the vaccine used for the primary course of vaccination against CVI.

44,702 people were revaccinated in Kazakhstan. In total, more than 8.8 million people (47.2% of the population) were vaccinated with the first component, more than 8.2 million people (43.3% of the population) were vaccinated with the second component.



WEST DALA LLP

**strongly recommends all employees
to get another vaccination against**

COVID-19

**in order to increase the protection
of their body from infection**



REVACCINATION AGAINST COVID-19

Revaccination of the population against coronavirus infection is conducted in Kazakhstan.

Taking into account the current epidemiological situation in the world, the National Advisory Commission on immunization has revised the tactics of revaccination of the population in the country since

December 8, 2021.



WHAT IS REVACCINATION FOR?

Over time, the level of protection of the body from infection decreases. The vaccination is repeated to activate the components of immunity, such as antibodies, B lymphocytes, T lymphocytes and T cells.

WHO IS BEING REVACCINATED?

- Revaccination against CVI is subject to all persons, including foreign citizens residing in Kazakhstan for three or more months, who have received a complete completed course of primary vaccination against CVI.
- Revaccination against COVID-19 is conducted once at intervals of six months after completion of the course of primary vaccination against COVID.
- Revaccination of pregnant women against CVI, who previously received the primary course of vaccination against CVI, is carried out using the Comirnaty vaccine (Pfizer, USA) in the period from the 16th to the 37th week of pregnancy.
- Revaccination against CVI of persons who have had CVI with a full course of primary vaccination against CVI is carried out no earlier than 6 months after recovery.



Also, by the resolution of the advisory commission on immunization, information on revaccination against CVI of Kazakhstanis who received revaccination against CVI abroad, when providing supporting documents is planned to be included in the module "Vaccination" of the Register of population enlistment of the Ministry of Healthcare of the Republic of Kazakhstan.